Applicant: Ilya

Ilya Trakht et al.

Serial No.:

09/664,958

Filed: Page 9

September 18, 2000

Amendments to the Abstract:

Please replace the abstract with the abstract amended as indicated below, with deleted matter indicated by strikethrough and added matter indicated by underlining. A clean version of

the amended abstract is attached hereto as Exhibit A.

The present This invention provides monoclonal antibodyproducing hybridomas designated 27.F7 and 27.B1. invention also provides a method of methods for detecting TIP-2 antigen bearing antigen-bearing cancer cells in a sample. The invention provides a method of , detecting the presence of TIP-2 antigen, optionally on the surface of cells. The invention provides a method immunohistochemical screening of a tissue section for the presence of TIP-2 antiqen bearing cancer cells, diagnosing cancer in a subject. The invention provides a method , monitoring progression of cancer wherein the cancer cells are TIP-2 antigen-bearing cells, delivering exogenous material to TIP-2 antigen-bearing cancer cells of a human subject. The invention provides a method , and treating cancer in a human The This invention further provides a kit for subject. detecting the presence of TIP-2 antigen-bearing cancer cells. This invention also provides isolated peptides having the amino acid sequences Lys Leu Leu Gly Gly Gln Ile Gly Leu (SEQ. ID No.) (SEQ ID NO:3) and Ser Leu Leu Gly Cys Arg His Tyr Glu Val (SEQ. ID No.) <u>(SEQ ID NO:4)</u>. The invention provides a method for immunohistochemical screening of a tissue section for the presence of TIP-2 antigen bearing cancer cells. The invention provides a kit for detecting the presence of TIP-2 antigen-bearing cancer cells. The invention provides a method for detecting the presence of TIP-2 antigen. The invention provides a method for

 \int_{0}^{∞}

Applicant :

Ilya Trakht et al.

Serial No.:

09/664,958

Filed:

September 18, 2000

Page 10

immunohistochemical screening of tissue sections. The invention provides a method for monitoring progression of cancer wherein the cancer cells are TIP-2 antigen-bearing cells. The invention provides a method for diagnosing cancer associated with the expression of TIP-2.